

Comment on "Theory of Unconventional Spin Density Wave: A Possible Mechanism of the Micromagnetism in U-based Heavy Fermion Compounds"

In the recent letter [1] a new, very attractive idea is proposed for the explanation of the micromagnetism in U-based heavy fermion (HF) compounds. For this sake a nontrivial spin density wave (SDW) state is introduced in the framework of the Hamiltonian:

$$H = -t \sum_{\langle ij \rangle \sigma} (c_{i\sigma}^\dagger c_{j\sigma} + h.c.) + U \sum_i n_{i\uparrow} n_{i\downarrow} - 2J \sum_{\langle ij \rangle} \vec{S}_i \vec{S}_j + (V - \frac{J}{2}) \sum_{\langle ij \rangle \sigma \sigma'} n_{i,\sigma} n_{j,\sigma'} \quad (1)$$

where we used the same notations as in [1] except $\vec{S}_i = \frac{1}{2} c_i^\dagger \vec{\sigma} c_i$. Unlike the conventional SDW, the order parameter $\Psi_k^Q \equiv \sum_{\sigma} \sigma \langle c_{k\sigma}^\dagger c_{k+Q\sigma} \rangle$ in the unconventional SDW (d-SDW) [1] state is characterized by "d-wave" like k-dependence $\Psi_k^Q \propto \cos k_x - \cos k_y$ [2]. In this case the ordered staggered magnetic moment M_Q is equal to zero. The authors restricted themselves to a very special case of 2D electron system on a simple square lattice, the shape of the Fermi surface corresponding to the *perfect nesting* with $Q=(\pi, \pi)$. The direct and exchange interaction constants are chosen positive $V > 0$, $J > 0$ [3].

We are not going to discuss the origin of the model (1) and criticize its applicability to the essentially 3D HF compounds (such as UPt_3 and URu_2Si_2 [4]) without any experimental evidence of perfect or imperfect nesting. Our goal is to claim, that even in the model considered in [1], the mean field (MF) analysis performed by the authors is incomplete and the phase diagram obtained (see Fig.1 in [1]) is wrong.

To begin with, let us look carefully on the Hamiltonian (1). One can easily see, that this Hamiltonian contains the Coulomb interaction and the *ferromagnetic (!)* [3] exchange integral. Thus, there are at least four ordered states which may be realized in this model: itinerant ferromagnet (FM) state, conventional SDW, charge density wave (CDW), and d-SDW. One can expect, that the FM state, missed by [1], will be dominant at least in the limit $U, V \ll J$. Therefore, to construct a complete phase diagram, the FM state should also be incorporated into the MF approach.

Let us consider first the case $(U, V, J) \ll t$ when the nesting property is important and MF analysis is reasonable. The criterion of instability can be determined from the behavior of the static response functions [5]: $\chi_\alpha(q, 0) = \chi_\alpha^0(q, 0) / (1 - I_\alpha(q) \chi_\alpha^0(q, 0))$, where $\alpha = FM, DW$, $I_{FM}(0) = U + 4J$, $I_{SDW}(Q) = U - 4J$, $I_{CDW}(Q) = 8V - U - 4J$ and $I_{d-SDW}(Q) = V$. For the perfect nesting case $\chi_{DW}^0(Q, 0) \sim (1/t) \log^2(t/T)$ [6], where one power of logarithm comes from nesting and

another one is due to the Van Hove singularity (VHS). Nevertheless, $\chi_{FM}^0(0, 0) \sim (1/t) \log(t/T)$ is also singular [6] due to VHS. The MF critical temperatures [7] are $T_{DW}^{MF} \sim t \exp(-2\pi\lambda_{DW}\sqrt{t/I_{DW}})$ and $T_{FM}^{MF} \sim t \exp(-2\pi\lambda_{FM}t/I_{FM})$, $\lambda_\alpha \sim 1$. Thus, the FM state certainly wins when $J \gg (U, V)$ and $V/J \lesssim J/t$ and even overcomes d-SDW in the phase diagram Fig.1 in [1]. The SDW state is more favorable when $U \gg (V, J)$ and CDW state occurs when $V \gg (U, J)$. We also emphasize, that unlike VHS, an additional "nesting" singularity in $\chi(Q, 0)$ is very sensible to a variety of effects, such as interlayer tunneling, doping, next hopping, etc, making the application of model [1] to real systems nearly impossible.

Let us consider another important limit $U \gg (t, V, J)$, the most realistic one, since the one-site U should be larger than the other nearest-neighbor interactions V, J and $t \sim m_*^{-1} \ll t_0$ ($m_* \gg m_0$ is an effective HF mass, m_0 and t_0 correspond to noninteracting fermions). In this case the V term is irrelevant for the half-filled band due to the constraint $n_i=1$, nesting is not important and only the AF state with $I_{AF} \sim t^2/U$ [8] is possible (when $J/t < t/U$).

To conclude, the new d-SDW state predicted in [1] cannot be realized for the most physically reasonable limits. The phase diagram in [1] is wrong, resulting in an erroneous statement of the d-SDW stability region. The very narrow region of parameters U, V, J, t (which has nothing to do with those presented in [1]) where the d-SDW state may exist requires a more detailed analysis.

We would like to thank K. Kikoin, F. Onufrieva and P. Pfeuty for discussions. M.K. is grateful for the support and hospitality of Laboratoire Léon Brillouin (CE-Saclay) where this work was carried out.

M.N. Kiselev and F. Bouis

Laboratoire Léon Brillouin, CE-Saclay 91191 Gif-sur-Yvette Cedex, France

-
- [1] H. Ikeda and Y. Ohashi. Phys. Rev. Lett. **81**, 3723 (1998).
 - [2] d-SDW is more favorable among all other SDW (extended s- and p- states) due to the Van Hove singularity.
 - [3] There is not any unconventional SDW state for $J \leq 0$.
 - [4] N. Grewe and F. Steglich, in *Handbook of the Physics and Chemistry of Rare Earths*, edited by K.A. Gschneider Jr. and L. Eyring (Elsevier, Amsterdam, 1991), Vol. 14, p.343.
 - [5] For simplicity we use the random phase approximation.
 - [6] J.E. Hirsch, Phys. Rev. B **31**, 4403 (1985).
 - [7] There is no true transition to a state with long range order in a 2D system with a continuous symmetry except for $T=0$.
 - [8] P.W. Anderson. Phys. Rev. **115**, 2 (1959).